



## Evaluation of antioxidant enzymes activity in canola under salt stress

JafarAbili<sup>1</sup>, Sajjad Zare<sup>2\*</sup>

<sup>1</sup>Islamic Azad University, Abhar Branch, Abhar, Iran

<sup>2</sup> Agricultural Biotechnology Research Institute of Iran, PO Box 31535-1897, Karaj, Iran

\*zbreeder@gmail.com

### ABSTRACT

Salinity is one of the major stresses in arid and semi-arid regions causing adverse effects at physiological, biochemical, and molecular levels, limiting crop productivity. In this research, three canola cultivars (Licord, Talayeh, Zarfam) were compared at 5 salinity levels (control, 50, 100, 150 and 200 mM) for their catalase, guaiacol peroxidase, superoxide dismutase activity, proline and yield in a completely randomized design with 3 replications. In our study, we found that NaCl concentrations greater than 150 and 200 mM caused the irreversible disorders. Increased salt concentrations led to significant changes in the levels of antioxidative enzymes and proline in three canola cultivars. Also, yield rates in three varieties decreased in the presence of NaCl concentrations.

**KEY WORDS:** Catalase; Peroxidase; Stress; superoxide dismutase.



## Council for Innovative Research

Peer Review Research Publishing System

.Journal: JOURNAL OF ADVANCES IN AGRICULTURE

Vol 2., No 2

[jaaeditor@gmail.com](mailto:jaaeditor@gmail.com)

[www.cirjaa.com](http://www.cirjaa.com)

## INTRODUCTION

Salinity is one of the major stresses in arid and semi-arid regions causing adverse effects at physiological, biochemical, and molecular levels, limiting crop productivity, also Salinity and drought are most important problems in Iran's agriculture. Salinity induces water deficit even in well watered soils by decreasing the osmotic potential of soil solutes, thus making it difficult for roots to extract water from their surrounding media (Sairam et al., 2002). Although, Salinity contributes to formation of reactive oxygen species (ROS), superoxideradical ( $O_2^-$ ), hydrogen peroxide ( $H_2O_2$ ) and hydroxyl radical ( $OH^-$ ) but plants increase the activity of antioxidant enzymes for decreasing of salinity effects (Moller et al., 2007). Previously, this fact reported by some researcher such as Zare and Pakniyat (2012), Ashraf and Ali (2007). The capability of scavenging ROS and reducing their damaging effects may correlate with the salinity tolerance of plants (Guo et al., 2006), therefore, At this article, we studied yield and antioxidant enzymes activity of three canola cultivars to salinity stress.

## MATERIAL AND METHODS

Three canola cultivars (Licord, Talayeh, Zarfam) were compared at 5 salinity levels (control, 50, 100, 150 and 200mM) for their catalase, guaiacol peroxidase, superoxide dismutase activity and proline in a completely randomized design with 3 replications. Before experimentation, all the seed samples were surface sterilized with 10% sodium hypochlorite solution for 5 min and washed three times with sterilized distilled water. Salt was applied to appropriate pots in split and in 5 stages within 5 weeks to final concentrations by irrigation based on soil field capacity, after 7 days of final salt treatment leave samples were collected and frozen in liquid nitrogen immediately and stored at  $-20^\circ C$  before analysis. Yield per pots were determined at end of experiment. The protein content was estimated according to Bradford (1976), using bovine serum albumin as a standard. Superoxide dismutase activity, the basis of which is its ability to inhibit the photochemical reduction of nitrobluetetrazolium (NBT) (Beauchamp and Fridovich, 1971), was determined according to the method of Dhindsa et al. (1980). For SOD assay, the reaction mixture contained 50mM K-phosphate buffer (pH 7.8), 13mM methionine, 75 $\mu M$  NBT, 0.1 $\mu M$  EDTA, 4 $\mu M$  riboflavin and required amount of enzyme extract. The reaction was started by adding riboflavin and placing the tubes under two 15 W fluorescent lamps for 15 min. A complete reaction mixture without enzyme, which gave the maximal colour, served as control. A non-irradiated complete reaction mixture served as a blank. One unit of SOD activity was defined as the amount of enzyme required to cause 50% inhibition of the reduction of NBT as monitored at 560 nm, which was measured according to the method of Giannopolitis and Ries (1977). Peroxidase activity was assayed adopting the method of Polle et al. (1994). According to this method POD activity was determined at 436 nm by its ability to convert guaiacol to tetraguaiacol ( $\epsilon = 26.6 \text{ mM}^{-1} \text{ cm}^{-1}$ ) The reaction mixture contained 100mM K-phosphate buffer (pH 7.0), 20.1mM guaiacol, 10mM  $H_2O_2$  and enzyme extract. The increase in absorbance was recorded by the addition of  $H_2O_2$  at 436 nm for 5 min. CAT activity was determined by monitoring the disappearance of  $H_2O_2$  at 240 nm ( $\epsilon = 40 \text{ mM}^{-1} \text{ cm}^{-1}$ ) according to the method of Aebi (1984). The reaction mixture contained 50mM K-phosphate buffer (pH 7.0), 33mM  $H_2O_2$  and enzyme extract.

## RESULTS AND DISCUSSION

**Catalase Activity:** according to the results, catalase activity decreased with increasing in salinity stress, these were observed in Licord, Talayeh and Zarfam under 200 in comparison with the control (43, 40 and 22%, respectively) (Fig 1). The decline in CAT activity is regarded as a general response to many stresses (Herbinger et al. 2002; Bakalova et al. 2004; Jung 2004; Guo et al. 2006; Pan et al. 2006; Gunes et al. 2008; Liu et al. 2008; Abedi and Pakniyat 2010; Zare and Pakniyat 2012). The reduction of CAT activity is supposedly due to the inhibition of enzyme synthesis or change in the assembly of enzyme subunits under stress conditions. It may also be associated with degradation caused by induced peroxisomal proteases or may be due to the photo-inactivation of the enzyme (Abedi and Pakniyat 2010; Zare and Pakniyat 2012). In normal condition, Licord had the highest catalase activity but this decrease was slowly in Zarfam cultivar.

**Peroxidase Activity:** A significant increase ( $P < 0.01$ ) was observed in peroxidase activity under stress especially 200mM for all three cultivars Licord, Talayeh and Zarfam (10, 8.1 and 9.23%, respectively) when compared to the control (Fig 2). The increase in expression of guaiacol peroxidase could be part of the oilseed rape's response to the oxidative damage caused by the increasing levels of salt stress (Zare and Pakniyat 2012). Some previous studies, as parallel with our results, mentioned the increased POD activity under salinity stress conditions in other plants such as pea (Hernandez et al., 2000) rice (Lee et al., 2001) wheat (Sairam et al., 2002).

**Superoxide Dismutase Activity:** Highest activity was observed at 100mM treatment but activity of this enzyme reduced by 150 and 200 mM treatments (Fig 3). This may be related to the low potential of canola plants to remove  $O_2^-$  under high concentration of salt. Cultivars were similar regarding their superoxidase activity at all salt levels. According to this fact that SOD processing is known to be substrate inducible (Tsang et al. 1991), an increase in the SOD activity may be attributed to the increased production of active oxygen species as substrate that lead to increased expression of genes encoding SOD (Abedi and Pakniyat 2010).

**Proline Content:** At all NaCl concentrations, Proline content recorded in three varieties increased significantly compared to control. For plants grown in the presence of 200 mM NaCl, the proline content in Licord, Talayeh, Zarfam increased 1.52, 1.55 and 1.61 times compared to control plants, respectively (Fig 4). Proline is known to accumulate under drought and saline conditions (Misra and Gupta, 2005; Jaleel et al., 2007; Yang and Lan, 2009).

**Yield:** According to results, yield rates in three varieties decreased in the presence of NaCl concentrations. In the presence of 200 mM NaCl, the yield in Licord, Talayeh, Zarfam decreased 85, 89 and 91% compared to control plants, respectively, (Fig 5). As a result, the induction of antioxidant enzyme activities is a general adaptation strategy which



plants use to overcome oxidative stresses (Foyer & Noctor 2003). In our study, we found that NaCl concentrations greater than 150 and 200 mM caused the irreversible disorders. In this study, increased salt concentrations led to significant changes in the levels of antioxidative enzymes in three canola cultivars. Peroxidase and superoxidase increased by salinity stress but catalase activity decreased. Similar responses have been observed in *B. maritima* and *B. vulgaris* (Bor et al., 2003), cotton (Meloni et al., 2003), maize (Neto et al., 2006), and cabbage (Posmyk et al., 2009). We found that the 200mM treatment had a higher proline concentration, a positive correlation between abiotic stress tolerance and free proline accumulation has been reported (Martinez et al., 2003) Also, our result demonstrated that Licord, Talayeh, Zarfam had similar response patterns to salinity.




Fig 1: Effect of NaCl treatments on CAT activity in leaves of three cultivars of *Brassica napus* (Means  $\pm$ SE)




Fig 2: Effect of NaCl treatments on Peroxidase activity in leaves of three cultivars of *Brassica napus* (Means  $\pm$ SE)




Fig 3: Effect of NaCl treatments on Superoxide dismutase activity in leaves of three cultivars of *Brassica napus* (Means  $\pm$ SE)




Fig 4: Effect of NaCl treatments on Proline content in leaves of three cultivars of *Brassica napus* (Means  $\pm$ SE)




Fig 5: Effect of NaCl treatments on Yield of three cultivars of *Brassica napus* (Means  $\pm$ SE)



## References

1. Abedi, T., Pakniyat, H. 2010. Antioxidant Enzyme Changes in Response to Drought Stress in Ten Cultivars of Oilseed Rape (*Brassica napus* L.). *Czech J. Genet. Plant Breed.*, 46, 2010 (1): 27–34
2. Ashraf, M., and Q. Ali, 2008. Relative membrane permeability and activities of some antioxidant enzymes as the key determinants of salt tolerance in canola (*Brassica napus* L.). *Environ. Exp. Bot.*, 63: 266-273.
3. Bakalova S., Nikolova A., Wedera D. 2004. Isoenzyme profiles of peroxidase catalase and superoxide dismutase as affected by dehydration stress and ABA during germination of wheat seeds. *Journal of Plant Physiology*, 30: 64–77.
4. Bor M, Özdemir F, Türkan I. 2003. The effect of salt stress on lipid peroxidation and antioxidants in leaves of sugar beet *Beta vulgaris* L. and wild beet *Beta maritima* L. *Plant Soil* 164: 77- 84,
5. Foyer C., Noctor G. 2003. Redox sensing and signaling associated with reactive oxygen in chloroplasts, peroxisomes and mitochondria. *Physiologia Plantarum*, 119: 355–364.
6. Gunes A., Pilbeam D., Inal A., Coban S. 2008. Influence of silicon on sunflower cultivars under drought stress, I: Growth, antioxidant mechanisms and lipid peroxidation. *Commun. Soil Science & Plant Nutrition*, 39: 1885–1903.
7. Guo Z., Ou W., Lu S., Zhong Q. 2006. Differential responses of antioxidative system to chilling and drought in four rice cultivars differing in sensitivity. *Plant Physiology and Biochemistry*, 44: 828–836.
8. Herbinger K., Tausz M., Wonisch A., Soja G., Sorger A., Grill D. 2002. Complex interactive effects of drought and ozone stress on the antioxidant defence systems of two wheat cultivars. *Plant Physiology and Biochemistry*, 40: 691–696.
9. Hernandez, J. A., A. Jimenez, P. Mullineaux, F. Sevilla, 2000. Tolerance of pea (*Pisumsativum* L.) to long-term salt stress is associated with induction of antioxidant defences. *Plant Cell Environ.*, 23: 853-862.
10. Jaleel CA, Gopi R, Sankar B. 2007. Studies on germination, seedling vigour, lipid peroxidation, and proline metabolism in *Catharanthus roseus* seedlings under salt stress. *S Afr J Bot* 73: 190-195,
11. Jung S. 2004. Variation in antioxidant metabolism of young and mature leaves of *Arabidopsis thaliana* subjected to drought. *Plant Science*, 166: 459–466.
12. Lee, D. H., Y. S. Kim, C. B. Lee, 2001. The inductive responses of the antioxidant enzymes by salt stress in the rice (*Oryza sativa* L.). *J. Plant Physiol.*, 158: 737-745
13. Liu J., Xie X., Du J., Sun J., Bai X. 2008. Effects of simultaneous drought and heat stress on Kentucky bluegrass. *Journal of Horticultural Science*, 115: 190–195.
14. Martinez CA, Maestri M, Lani EG. 2003. In vitro salt tolerance and proline accumulation in Andean potato (*Solanum* spp.) differing in frost resistance. *Plant Sci* 116: 117-184,
15. Meloni DA, Oliva MA, Martinez CA. 2003. Photosynthesis and activity of superoxide dismutase peroxidase and glutathione reductase in cotton under salt stress. *Environ Exp Bot* 49: 69- 76,.
16. Misra N, Gupta AK. 2005. Effect of salt stress on proline metabolism in two high yielding genotypes of green gram. *Plant Sci* 169: 331-339.
17. Moller, I. M., P. E. Jensen, A. Hansson, 2007. Oxidative modifications to cellular components in plants. *Ann. Rev. Plant Biol.*, 58: 459-481.
18. Neto ADA, Prisco JT, Enéas-Filho J. 2006. Effect of salt stress on antioxidative enzymes and lipid peroxidation in leaves and roots of salt-tolerant and salt-sensitive maize genotypes. *Environ Exp Bot* 56: 87-94,
19. Pan Y., Wu L.J., Yu Z.L. 2006. Effect of salt and drought stress on antioxidant enzymes activities and SOD isoenzymes of liquorice (*Glycyrrhizauralensis* Fisch). *Journal of Plant Growth Regulation*, 49: 157–165.
20. Posmyk MM, Kontek R, Janas KM. 2009. Antioxidant enzymes activity and phenolic compounds content in red cabbage seedlings exposed to copper stress. *Ecotox Environ Safe* 72: 596-602,
21. Sairam, R. K., K. V. Rao, and G. C. Srivastava, 2002. Differential response of wheat genotypes to long-term salinity stress in relation to oxidative stress, antioxidant activity and osmolyte concentration. *Plant Sci.*, 163: 1037-1046.
22. Tsang E.W.T., Bowler C., Herouart D., Van C.W., Villarroel R., Genetello C., Van M.M., Inze D. 1991. Differential regulation of superoxide dismutase in plants exposed to environmental stress. *Plant Cell*, 3: 783–792.
23. Yang SL, Lan SS, 2009. Gong M. Hydrogen peroxide-induced proline and metabolic pathway of its accumulation in maize seedlings. *J Plant Physiol* 166: 1694-1699.
24. Zare, S., Pakniyat, H. 2012. Changes in activities of antioxidant enzymes in oilseed rape in response to salinity stress. *International Journal of Agriculture and Crop Sciences*. 4-7/398-403